

Development of QED-cascade arising at reflection of a PW laser pulse from a solid nanostructured target

Serebryakov M.A.^{1,®}, Nerush E.N.¹, Kostyukov I.Yu.¹,
Serebryakov M.A.¹, Nerush E.N.¹ and Kostyukov I.Yu.¹

¹ Institute of Applied Physics of the Russian Academy of Sciences, Ulyanova 46, Nizhny Novgorod, 603950, None

® serebryakovma@ipfran.ru

Quantum electrodynamics (QED) cascade arising in a superposition of incident multipetawatt laser pulse and its reflection from a solid target were studied. It is numerically [1] shown that with normalized laser field amplitude $a_0 = eE_0/(mc\omega) = 250$, pulse duration 120 fs and density $n_e = 500n_{cr}$ (which approximately corresponds, for example, to a diamond target; $n_{cr} = m\omega^2/[4\pi e^2]$), it is possible to obtain QED cascade in this setup. The growth rate of the number of particles is very close to that in ideal linear-polarized standing wave [2]. Consequently, it is not necessary to use multiple laser channels to observe a QED cascade as it was suggested earlier [3, 4]. One laser channel and a reflective plasma target are sufficient, which can make the experiment more simple for a lack of necessity of operating with multiple laser pulses and its precision synchronization.

[1] QUILL, <https://github.com/QUILL-PIC/Quill>

[2] Grismayer T, Vranic M, Martins J L, Fonseca R A and Silva L O 2017 *Physical Review E* **95** 023210

[3] Gonoskov A, Bashinov A, Bastrakov S, Efimenko E, Ilderton A, Kim A, Marklund M, Meyerov I, Muraviev A and Sergeev A 2017 *Physical Review X* **7** 041003 ISSN 2160-3308

[4] Nerush E N, Iligenov R R and Kostyukov I Y 2023 *Bulletin of the Lebedev Physics Institute* **50** S689–S692 ISSN 1934-838X