

Anomalies of spatial ions distribution in trap with different restraining potential

Chigvintsev A Yu^{1,®}, Iosilevskiy I L^{1,2}, Noginova I Yu^{3,4} and Zorina I G⁵

¹ Moscow Institute of Physics and Technology, Institutskiy Pereulok 9, Dolgoprudny, Moscow Region 141701, Russia

² Joint Institute for High Temperatures of the Russian Academy of Sciences, Izhorskaya 13 Bldg 2, Moscow 125412, Russia

³ Russian State Agrarian University—Timiryazev Moscow Agricultural Academy, Timiryazevskaya 49, Moscow 127550, Russia

⁴ National University of Science and Technology “MISIS”, Leninskiy Avenue 4, Moscow 119049, Russia

⁵ Bauman Moscow State Technical University, 2nd Baumanskaya Street 5, Moscow 105005, Russia

® alex012008@gmail.com

Discontinuities in equilibrium spatial charge profiles of ions in non-uniform Coulomb systems is a common phenomenon in wide number of thermoelectrostatics problems. As we showed previously, such discontinuities are peculiar micro-level manifestation of phase transitions and intrinsic macro-level non-ideality elects in local equation of state (EOS), which should be used for description of non-ideal ionic subsystem in frames of local-density (or “pseudo fluid”, or “jellium” etc) approximation. Special emphasis is made in present paper on the mentioned above non-ideality elects in non-uniform ionic subsystems, such as equilibrium charge profile in ionic traps with different “external potentials”. Multiphase EOS for simplified ionic model of classical charged hard spheres on uniformly compressible electrostatic compensating background was constructed and several illustrative examples of discussed discontinuous ionic profiles were calculated.