

# Jones–Wilkins–Lee equation of state for high-explosion detonation products based on cylinder-test experiments using synchrotron diagnostics

**Smirnov E B<sup>1,®</sup>, Petrov D V<sup>1</sup>, Garmashev A Yu<sup>1</sup>, Muzyrya A K<sup>1</sup>, Prosvirnin K M<sup>1</sup>, Shakhmaev S V<sup>1</sup>, Puuel E R<sup>2</sup>, Ten K A<sup>2</sup> and Kashakarov A O<sup>2</sup>**

<sup>1</sup> Federal State Unitary Enterprise “Russian Federal Nuclear Center—All-Russian Research Institute of Experimental Physics”, Mira Avenue 37, Sarov, Nizhny Novgorod Region 607188, Russia

<sup>2</sup> Lavrentyev Institute of Hydrodynamics of the Siberian Branch of the Russian Academy of Sciences, Lavrentyev Avenue 15, Novosibirsk 630090, Russia

® ewgeny\_smirnov@mail.ru

The work by detonation products (DP) on the environment in a process and shock-wave parameters generated within the environment for each high explosive (HE) are definitely defined by the equation of state (EOS) for DP. The DP EOS is generally determined based on the DP isentrope passing through the Chapman–Jouguet point. The experiments to determine the HE throwing action are conducted to establish the behavior of the DP expansion isentrope. For this purpose, common are the experiments on end-face throwing of plates under the effect of HE DP and the experiments on throwing cylindrical shells, so-called cylinder-test experiments. The presence of the metal shell in the experiment contributes to distortion of the HE DP expansion process but remains necessary to conduct optical and electric-contact diagnostics. This work suggests the revised version of the cylinder-test experiments without metal shell involving diagnostics of DP expansion through synchrotronic radiation. The experiments were conducted using the accelerator complex VEPP-3 of the Budker Institute of Nuclear Physics of the Siberian Branch RAS. The experimental data were used to plot isentropic expansion of the HE DP for the Jones–Wilkins–Lee EOS.