

ANOMALOUS SPATIAL CHARGE PROFILES OF PLASMA AS MANIFESTATION OF PHASE TRANSITIONS IN MODIFIED ONE COMPONENT PLASMA MODEL

*Chigvintsev A. Yu.,^{*1} Iosilevskiy I.L.,^{1,2} Noginova L. Yu.,³ Zorina I. G.⁴*

¹MIPT, Dolgoprudny, Russia, ²JIHT RAS, Moscow, Russia, ³MISiS, Moscow, Russia, ⁴BMSTU, Moscow, Russia

^{*}alex012008@gmail.com

The paper discusses the possibility of the appearance of discontinuities in the results of calculations of equilibrium space charge profiles in the vicinity of the source of inhomogeneity [1]. These discontinuities are considered as a kind of micro-level manifestation of phase transitions and other (macro-level) charge correlation effects (“non-ideality”) contained in the local equation of state (EOS), which is used to describe the non-ideal electronic and/or ionic subsystem within the framework of the quasi-homogeneity approximation (“local density”) [2]. Particular attention in this work is paid to the possibility of a specific manifestation of the above-mentioned nonideality effects in the studied equilibrium charge profiles in the form of an ultradisperse two-phase mixture (“mixed phase”). The proposed general conclusion is the statement that the concept of mixed phase is not an attribute of exclusively astrophysical applications, but is a fairly general property of computational schemes used to describe equilibrium inhomogeneous Coulomb systems [3].

-
1. Iosilevskiy I. L., Chigvintsev A. Yu., Noginova L. Yu., Zorina I. G. // High Temperature, 60, Suppl. 3, S325-S331 (2022)
 2. Iosilevski I. L. // High Temperature 1985 V. 23 P. 807
 3. Chigvintsev A. Yu., Iosilevskiy I. L., Noginova L. Yu., Zorina I. G. J. of Phys. Conf. Ser. 2018. V. 946 012092