

Structures of active Brownian particles in a monolayer of colloidal plasma under laser irradiation

Zamorin D.A.^{1,2,@}, **Syrovatka R.A.**¹, **Koss X.G.**^{1,2},
Vasiliev M.M.¹ and **Petrov O.F.**¹

¹ Joint Institute for High Temperatures of the Russian Academy of Sciences,
Izhorskaya 13 Bldg 2, Moscow, 125412, Russia

² Moscow Institute of Physics and Technology, Institutskiy Pereulok 9,
Dolgoprudny, Moscow Region, 141701, Russia

@ zamorin.da@phystech.edu

This work presents experimental results on structural transitions in a monolayer of active Brownian particles. The experiments were conducted using synthesized melamine-formaldehyde (MF) particles confined in a radio-frequency (RF) plasma discharge. The particles were coated with a light-absorbing layer via magnetron sputtering, enabling localized heating under laser irradiation and thereby increasing their kinetic energy of motion. A transition from an ordered to a disordered state was observed in a system far from thermodynamic equilibrium, proceeding through an intermediate hexatic state [1]. The analysis of this transition employed pair correlation functions, bond-angular correlation functions, the two-dimensional static structure factor, and a characterization of topological defects. The findings indicate a two-stage melting scenario consistent with the Berezinskii–Kosterlitz–Thouless (BKT) theory [2] for two-dimensional systems in thermodynamic equilibrium. This work was supported by the Russian Science Foundation (Project No. 25-12-00406).

[1] Vasilieva E V P O F and M V M 2021 *Sci Rep.* **11**(523)

[2] M K J and J T D 1973 *J. Phys. C: Solid State Phys.* **6**